



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: January 9, 2017
ACCEPTED: February 27, 2017
PUBLISHED: March 14, 2017

Analytic treatment of the excited instability spectra of the magnetically charged SU(2) Reissner-Nordström black holes

Shahar Hod

*The Ruppin Academic Center,
Emeq Hefer 40250, Israel*

*The Hadassah Academic College,
Jerusalem 91010, Israel*

E-mail: shaharhod@gmail.com

ABSTRACT: The magnetically charged SU(2) Reissner-Nordström black-hole solutions of the coupled nonlinear Einstein-Yang-Mills field equations are known to be characterized by infinite spectra of unstable (imaginary) resonances $\{\omega_n(r_+, r_-)\}_{n=0}^{\infty}$ (here r_{\pm} are the black-hole horizon radii). Based on direct *numerical* computations of the black-hole instability spectra, it has recently been observed that the excited instability eigenvalues of the magnetically charged black holes exhibit a simple universal behavior. In particular, it was shown that the numerically computed instability eigenvalues of the magnetically charged black holes are characterized by the small frequency universal relation $\omega_n(r_+ - r_-) = \lambda_n$, where $\{\lambda_n\}$ are dimensionless constants which are independent of the black-hole parameters. In the present paper we study analytically the instability spectra of the magnetically charged SU(2) Reissner-Nordström black holes. In particular, we provide a rigorous *analytical* proof for the *numerically*-suggested universal behavior $\omega_n(r_+ - r_-) = \lambda_n$ in the small frequency $\omega_n r_+ \ll (r_+ - r_-)/r_+$ regime. Interestingly, it is shown that the excited black-hole resonances are characterized by the simple universal relation $\omega_{n+1}/\omega_n = e^{-2\pi/\sqrt{3}}$. Finally, we confirm our analytical results for the black-hole instability spectra with numerical computations.

KEYWORDS: Black Holes, Classical Theories of Gravity

ARXIV EPRINT: [1701.01447](https://arxiv.org/abs/1701.01447)

Contents

1	Introduction	1
2	Description of the system	2
3	The characteristic resonance condition	4
4	The black-hole excited instability spectra	6
5	Numerical confirmation	7
6	Summary	8

1 Introduction

It is well known that the electrically charged U(1) Reissner-Nordström black-hole spacetime describes a *stable* solution of the coupled nonlinear Einstein-Maxwell field equations [1, 2] (see also [3–5]). On the other hand, the magnetically charged SU(2) Reissner-Nordström black-hole spacetime [6] describes an *unstable* solution of the coupled nonlinear Einstein-Yang-Mills field equations [7–13]. In fact, the Reissner-Nordström black-hole solutions of the Einstein-Yang-Mills theory are characterized by an *infinite* spectrum of unstable perturbation modes. These unstable (exponentially growing in time) modes are described by an infinite set of imaginary black-hole resonances $\{\omega_n\}_{n=0}^{n=\infty}$ [7–13].

In a very interesting numerical investigation of the coupled Einstein-Yang-Mills equations, it was recently revealed by Rinne [14] that these unstable magnetically charged black-hole solutions play the role of approximate¹ codimension-two intermediate attractors (critical solutions²) in the nonlinear gravitational collapse of the Yang-Mills field [14, 16–19].³ In particular, it has been shown explicitly [14] that the time spent in the vicinity of an unstable magnetically charged SU(2) Reissner-Nordström black-hole spacetime during a near-critical gravitational collapse of the nonlinear Yang-Mills field can be quantified by the characteristic scaling law⁴

$$\tau = \text{const} - \gamma \ln |p - p^*|. \quad (1.1)$$

¹As discussed in [14], the SU(2) Reissner-Nordström black-hole spacetime is only an approximate intermediate attractor of the nonlinear gravitational collapse of the Yang-Mills field because it is characterized by an *infinite* family of exponentially growing (unstable) perturbation modes.

²See [15] for an excellent review on the black-hole critical phenomena in nonlinear gravitational collapse.

³It is worth noting that this physically interesting fact refers to type I and Type III nonlinear critical behaviors in gravitational collapse, see [14, 16–19] for details.

⁴Here the quantity $|p - p^*|$ provides a measure in the physical parameter space for the distance of the initial field data from the threshold (critical) solution of the nonlinear Einstein-Yang-Mills theory [15].

It is interesting to note that the critical exponents in the scaling behavior (1.1), which characterizes the nonlinear near-critical gravitational collapse of the Yang-Mills field, are directly related to the imaginary eigenvalues which characterize the instability spectrum of the corresponding magnetically charged Reissner-Nordström black-hole spacetime [14]:

$$\gamma = 1/|\omega_{\text{instability}}|. \quad (1.2)$$

It is therefore physically interesting to investigate the characteristic instability (imaginary) resonance spectra $\{\omega_n(r_+, r_-)\}_{n=0}^{n=\infty}$ ⁵ of these magnetically charged black-hole solutions of the coupled nonlinear Einstein-Yang-Mills equations.

In his important numerical work, Rinne [14] has recently determined numerically the first three imaginary (unstable) resonant frequencies which characterize the magnetically charged SU(2) Reissner-Nordström black-hole spacetimes.⁶ Subsequently, in [20] we have analyzed the detailed numerical data provided by Rinne [14] and revealed the intriguing fact that, to a good degree of accuracy, the numerically computed [14] excited instability eigenvalues of the magnetically charged SU(2) Reissner-Nordström black holes are characterized by the remarkably simple *universal* behavior

$$\omega_n(r_+ - r_-) = \lambda_n \quad \text{for} \quad \omega_n r_+ \ll (r_+ - r_-)/r_+, \quad (1.3)$$

where $\{\lambda_n\}$ are dimensionless constants which seem to be *independent* of the black-hole parameters.

The main goal of the present paper is to determine *analytically* the characteristic instability spectra of the magnetically charged SU(2) Reissner-Nordström black-hole solutions of the coupled Einstein-Yang-Mills theory. In particular, in this paper we shall provide a rigorous analytical proof for the validity of the numerically suggested [14, 20] universal behavior (1.3) which characterizes the excited instability spectra of the SU(2) Reissner-Nordström black-hole spacetimes.

2 Description of the system

The SU(2) Reissner-Nordström black-hole spacetime of mass M and unit magnetic charge is characterized by the spherically-symmetric line element [6]

$$ds^2 = -\left(1 - \frac{2m}{r}\right)dt^2 + \left(1 - \frac{2m}{r}\right)^{-1}dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (2.1)$$

where the radially dependent mass function $m = m(r)$ is given by⁷

$$m(r) = M - \frac{1}{2r}. \quad (2.2)$$

⁵Here r_{\pm} are the black-hole horizon radii [see eq. (2.3) below].

⁶As noted above, the SU(2) Reissner-Nordström black-hole solutions of the coupled nonlinear Einstein-Yang-Mills field equations are characterized by *infinite* spectra $\{\omega_n(r_+, r_-)\}_{n=0}^{n=\infty}$ of imaginary (unstable) resonant frequencies [10, 11]. Reference [14] has provided, for the first time, detailed numerical results for the first three resonant frequencies which quantify the instability growth rates of these magnetically charged black-hole spacetimes.

⁷We shall use natural units in which $G = c = \hbar = 1$.

The radii of the black-hole (outer and inner) horizons are given by

$$r_{\pm} = M \pm \sqrt{M^2 - 1}. \quad (2.3)$$

As shown in [21], linearized perturbation modes $\xi(r)e^{-i\omega t}$ ⁸ of the magnetically charged black-hole spacetime are governed by the wave equation

$$\left[\frac{d^2}{dx^2} + \omega^2 - U(x) \right] \xi = 0, \quad (2.4)$$

where the radial coordinate $x = x(r)$ is defined by the differential relation⁹

$$dx/dr = [1 - 2m(r)/r]^{-1}. \quad (2.5)$$

The effective radial potential which governs the Schrödinger-like wave equation (2.4) is given by [21]

$$U[x(r)] = -\frac{1}{r^2} \left[1 - \frac{2m(r)}{r} \right]. \quad (2.6)$$

It is worth emphasizing the fact that the radial function $U(x)$ in eq. (2.4), which determines the spatial behavior of the black-hole perturbation modes, has the form of an effective binding potential. In particular, it is a negative definite function of the radial coordinate x and it vanishes asymptotically at the two boundaries $x \rightarrow \pm\infty$ of the magnetically charged black-hole spacetime. As shown in [21], well-behaved perturbation modes of the black-hole spacetime are characterized by spatially bounded (exponentially decaying) radial eigenfunctions at the two asymptotic boundaries:

$$\xi(x \rightarrow -\infty) \sim e^{|\omega|x} \rightarrow 0 \quad (2.7)$$

and

$$\xi(x \rightarrow \infty) \sim x e^{-|\omega|x} \rightarrow 0, \quad (2.8)$$

where $\omega = i|\omega|$.

The radial differential equation (2.4), supplemented by the physically motivated boundary conditions (2.7) and (2.8) [21], determine the discrete family $\{\omega_n(r_+, r_-)\}_{n=0}^{n=\infty}$ of unstable ($\Im\omega > 0$) resonances which characterize the SU(2) Reissner-Nordström black-hole spacetimes [9–11]. Interestingly, below we shall show explicitly that the characteristic resonance spectrum of the magnetically charged black holes can be studied analytically in the regime $|\omega_n|r_+ \ll 1$ of small imaginary resonant frequencies. In particular, we shall derive a remarkably compact *analytical* formula [see eq. (4.3) below] for the excited instability eigenvalues which characterize the SU(2) Reissner-Nordström black-hole solutions of the coupled Einstein-Yang-Mills theory.

⁸Note that unstable (exponentially growing in time) perturbation modes of the magnetically charged black-hole spacetime are characterized by the relation $\Im\omega > 0$.

⁹Note that the near-horizon region $r \rightarrow r_+$ of the black-hole spacetime is mapped by the differential relation (2.5) to $x \rightarrow -\infty$, whereas spatial infinity $r \rightarrow \infty$ is mapped to $x \rightarrow \infty$.

3 The characteristic resonance condition

The recent numerical results of Rinne [14] reveal that the excited instability resonances $\{\omega_n\}_{n=1}^{n=\infty}$ of the magnetically charged SU(2) Reissner-Nordström black-hole spacetimes are characterized by the property

$$|\omega_n|r_+ \ll 1; \quad n = 1, 2, 3, \dots \quad (3.1)$$

As we shall now show, the Schrödinger-like differential equation (2.4), which governs the dynamics of the black-hole perturbation modes, is amenable to an *analytical* treatment in the regime (3.1) of *small* resonant frequencies.

It was pointed out in [12, 13] that, in the $M \gg 1$ regime (the regime of weakly-magnetized SU(2) Reissner-Nordström black holes), the Schrödinger-like perturbation equation (2.4) can be transformed using the well-known Chandrasekhar transformations [22] to the physically equivalent Teukolsky-like radial equation [23]:

$$\Delta^2 \frac{d^2 \psi}{dr^2} + [\omega^2 r^4 + 2iM\omega r^2 - \Delta[2i\omega r + \ell(\ell+1)]]\psi = 0, \quad (3.2)$$

where the complex number

$$\ell \equiv \frac{-1 + i\sqrt{3}}{2} \quad (3.3)$$

plays the role of an effective spherical harmonic index (see [12, 13] for details), and [23]

$$\Delta(r; M \gg 1) = r^2 - 2Mr. \quad (3.4)$$

The mathematical Chandrasekhar transformations [22] can also be used in the case of generic magnetically charged SU(2) Reissner-Nordström black holes,¹⁰ in which case the generalized expression for the radial function $\Delta(r)$ in the Teukolsky-like radial perturbation equation (3.2) is given by [23]

$$\Delta(r; M) = r^2 - 2Mr + 1. \quad (3.5)$$

It is convenient to use the dimensionless physical variables [24–26]

$$z \equiv \frac{r - r_+}{r_+ - r_-}; \quad k \equiv -i\omega(r_+ - r_-); \quad \varpi \equiv -i\omega \frac{2Mr_+}{r_+ - r_-}, \quad (3.6)$$

in terms of which the radial differential equation (3.2) reads

$$z^2(z+1)^2 \frac{d^2 \psi}{dz^2} + [-k^2 z^4 + 2kz^3 - \ell(\ell+1)z(z+1) - \varpi(2z+1) - \varpi^2]\psi = 0. \quad (3.7)$$

¹⁰We use here the term ‘generic SU(2) Reissner-Nordström black holes’ to describe magnetically charged black holes with generic masses (that is, in the present analysis we do *not* assume the strong inequality $M \gg 1$ previously assumed in [12, 13]).

The physically acceptable solution¹¹ of the radial perturbation equation (3.7) in the near-horizon $kz \ll 1$ region¹² can be expressed in terms of the familiar hypergeometric function [24–27]:

$$\psi(z) = z^{1+\varpi}(z+1)^{1-\varpi} {}_2F_1(-\ell+1, \ell+2; 2+2\varpi; -z). \quad (3.8)$$

The physically acceptable solution¹³ of the radial perturbation equation (3.7) in the asymptotic region $z \gg \varpi+1$ ¹⁴ can be expressed in terms of the familiar confluent hypergeometric function [24–27]:

$$\psi(z) = Ae^{kz}z^{\ell+1} {}_1F_1(\ell+2; 2\ell+2; -2kz) + Be^{kz}z^{-\ell} {}_1F_1(-\ell+1; -2\ell; -2kz). \quad (3.9)$$

For small resonant frequencies in the regime (3.1), the values of the dimensionless coefficients A and B in (3.9) can be determined by matching the two solutions [(3.8) and (3.9)] of the radial perturbation equation (3.7) in the overlap region¹⁵

$$\varpi+1 \ll z \ll 1/k. \quad (3.10)$$

This matching procedure yields the expressions [24–26]

$$A = \frac{\Gamma(2\ell+1)\Gamma(2+2\varpi)}{\Gamma(\ell+2)\Gamma(\ell+1+2\varpi)} \quad (3.11)$$

and

$$B = \frac{\Gamma(-2\ell-1)\Gamma(2+2\varpi)}{\Gamma(-\ell+1)\Gamma(-\ell+2\varpi)} \quad (3.12)$$

for the dimensionless coefficients $\{A, B\}$ in (3.9).

Substituting the expressions (3.11) and (3.12) into the far-region expression (3.9) of the radial eigenfunction and using the asymptotic ($z \gg 1$) properties of the confluent hypergeometric functions [27], one finds the large- z asymptotic behavior [24–26]

$$\psi(z \rightarrow \infty) = \psi_1 r e^{-kz} + \psi_2 r^{-1} e^{kz} \quad (3.13)$$

of the radial eigenfunctions, where

$$(r_+ - r_-)\psi_1 = \frac{(2\ell+1)\Gamma^2(2\ell+1)\Gamma(2+2\varpi)}{\Gamma^2(\ell+2)\Gamma(\ell+1+2\varpi)}(-2k)^{-\ell} - \frac{(2\ell+1)\Gamma^2(-2\ell-1)\Gamma(2+2\varpi)}{\Gamma^2(-\ell+1)\Gamma(-\ell+2\varpi)}(-2k)^{\ell+1}, \quad (3.14)$$

¹¹That is, the mathematical solution which respects the physically motivated near-horizon boundary condition (2.7).

¹²Note that in the near-horizon region $kz \ll 1$ one can neglect the first two terms inside the square brackets in eq. (3.7) [24].

¹³That is, the mathematical solution which respects the physically motivated asymptotic boundary condition (2.8) at spatial infinity.

¹⁴Note that in the asymptotic region $z \gg \varpi+1$ one can neglect the last two terms inside the square brackets in eq. (3.7) [24].

¹⁵It is worth noting that the overlap radial region $\varpi \ll z \ll 1/k$ exists in the regime $|\omega|r_+ \ll 1$ of small resonant frequencies [see eqs. (3.1) and (3.6)].

and

$$(r_+ - r_-)^{-1} \psi_2 = \frac{(2\ell+1)\Gamma^2(2\ell+1)\Gamma(2+2\varpi)}{\ell(\ell+1)\Gamma^2(\ell)\Gamma(\ell+1+2\varpi)} (2k)^{-\ell-2} - \frac{(2\ell+1)(\ell+1)\Gamma^2(-2\ell-1)\Gamma(2+2\varpi)}{\ell\Gamma^2(-\ell)\Gamma(-\ell+2\varpi)} (2k)^{\ell-1}. \quad (3.15)$$

A spatially bounded (normalizable) radial eigenfunction which satisfies the physically motivated boundary condition (2.8) at spatial infinity [21] is characterized by the asymptotic relation $\psi(z \rightarrow \infty) \rightarrow 0$. Thus, the coefficient ψ_2 of the exploding exponent in the asymptotic expression (3.13) should vanish, yielding the characteristic resonance equation [see eq. (3.15)]

$$(2k)^{2\ell+1} = \left[\frac{\Gamma(2\ell+1)\Gamma(-\ell)}{(\ell+1)\Gamma(-2\ell-1)\Gamma(\ell)} \right]^2 \frac{\Gamma(-\ell+2\varpi)}{\Gamma(\ell+1+2\varpi)} \quad (3.16)$$

for the instability eigenvalues of the SU(2) Reissner-Nordström black-hole spacetimes. Taking cognizance of eq. (3.3), one can write the resonance equation (3.16) for the instability spectra of the magnetically charged black holes in the form¹⁶

$$k^{i\sqrt{3}} = 8^{i\sqrt{3}} e^{i2\pi/3} \frac{\Gamma^2(i\frac{\sqrt{3}}{2})}{\Gamma^2(-i\frac{\sqrt{3}}{2})} \frac{\Gamma(\frac{1-i\sqrt{3}}{2} + 2\varpi)}{\Gamma(\frac{1+i\sqrt{3}}{2} + 2\varpi)}. \quad (3.17)$$

4 The black-hole excited instability spectra

As we shall now show, the characteristic resonance equation (3.17) for the instability eigenvalues of the magnetically charged SU(2) Reissner-Nordström black holes can be solved analytically in the regime¹⁷

$$\varpi \ll 1. \quad (4.1)$$

In this small frequency regime the resonance equation (3.17) can be approximated by

$$k^{i\sqrt{3}} = 8^{i\sqrt{3}} e^{i2\pi/3} \frac{\Gamma^2(i\frac{\sqrt{3}}{2})}{\Gamma^2(-i\frac{\sqrt{3}}{2})} \frac{\Gamma(\frac{1-i\sqrt{3}}{2})}{\Gamma(\frac{1+i\sqrt{3}}{2})}, \quad (4.2)$$

which yields the characteristic *infinite* spectrum^{18,19}

$$\omega_n \times (r_+ - r_-) = i \times 8e^{-\frac{2\pi}{\sqrt{3}}(n-\frac{1}{3}) + \frac{4\theta-2\phi}{\sqrt{3}}}; \quad n = 1, 2, 3, \dots \quad (4.3)$$

of unstable ($\Im\omega > 0$) black-hole resonances, where

$$\theta \equiv \arg[\Gamma(i\sqrt{3}/2)]; \quad \phi \equiv \arg\left[\Gamma\left(\frac{1+i\sqrt{3}}{2}\right)\right]. \quad (4.4)$$

¹⁶Here we have used eq. (6.1.18) of [27].

¹⁷Note that this regime corresponds to $\omega r_+ \ll (r_+ - r_-)/r_+$ [see eq. (3.6)].

¹⁸Here we have used the relation $1 = e^{-i2\pi n}$, where the integer $n = 1, 2, 3, \dots$ is the resonance parameter of the black-hole perturbation mode. In addition, we have used here eq. (6.1.23) of [27].

¹⁹As noted in [12, 13], one finds the numerical ratio $\theta/2\phi = 1.0016$. Thus, one can replace, with an accuracy of 0.05%, the expression $(4\theta - 2\phi)/\sqrt{3}$ in (4.3) by the simpler term $\sqrt{3}\theta$.

r_+	10.0	8.0	6.0	4.0	2.0	1.5
$\omega_2^{\text{ana}}/\omega_2^{\text{num}}$	1.046	1.025	1.011	1.005	1.007	1.012

Table 1. The excited instability eigenvalues of the magnetically charged SU(2) Reissner-Nordström black holes. We display the dimensionless ratio $\omega_2^{\text{ana}}/\omega_2^{\text{num}}$, where $\{\omega_2^{\text{ana}}(r_+)\}$ are the analytically calculated instability eigenvalues of the SU(2) Reissner-Nordström black holes as given by the analytical formula (4.3) and $\{\omega_2^{\text{num}}(r_+)\}$ are the corresponding numerically computed [14] instability eigenvalues of the magnetically charged black holes. One finds a fairly good agreement between the numerical data of [14] and the analytically derived formula (4.3) for the excited instability spectra of the SU(2) Reissner-Nordström black holes.

It is worth emphasizing again that the analytically derived formula (4.3) for the characteristic instability spectra of the magnetically charged SU(2) Reissner-Nordström black holes is valid in the small frequency regime [see (3.6) and (4.1)]

$$\omega_n r_+ \ll \frac{r_+ - r_-}{r_+}. \quad (4.5)$$

This inequality implies that, for a given value of the black-hole dimensionless temperature $(r_+ - r_-)/r_+$, the analytical formula (4.3) describes an infinite family of unstable (imaginary) black-hole resonances in the regime

$$n \gtrsim \left\lceil \ln \left(\frac{r_+ - r_-}{r_+} \right) \right\rceil + 1. \quad (4.6)$$

It is interesting to note that the instability spectra (4.3) of the magnetically charged SU(2) Reissner-Nordström black-hole spacetimes have the simple generic form $\omega_n \times (r_+ - r_-) = \text{constant}_n \equiv \lambda_n$ [see eq. (1.3)]. We have therefore provided here an *analytical* proof for the *numerically*-observed [14, 20] universal behavior (1.3) of the black-hole excited instability eigenvalues.

5 Numerical confirmation

It is of considerable physical interest to verify the validity of the analytically derived formula (4.3) for the excited instability eigenvalues of the SU(2) Reissner-Nordström black holes. The corresponding instability eigenvalues of the magnetically charged black holes were recently computed numerically in the very interesting work of Rinne [14]. In table 1 we present the dimensionless ratio $\omega_2^{\text{ana}}/\omega_2^{\text{num}}$, where $\{\omega_2^{\text{ana}}(r_+)\}$ are the analytically calculated excited instability eigenvalues of the SU(2) Reissner-Nordström black holes as given by the analytical formula (4.3) and $\{\omega_2^{\text{num}}(r_+)\}$ are the numerically computed [14] instability eigenvalues of the magnetically charged black holes. From the data presented in table 1 one finds a fairly good agreement between the analytically derived formula (4.3) for the excited instability spectra of the SU(2) Reissner-Nordström black holes and the corresponding numerically computed black-hole instability eigenvalues.

6 Summary

The magnetically charged SU(2) Reissner-Nordström black-hole spacetimes describe a family of unstable solutions of the coupled nonlinear Einstein-Yang-Mills field equations [7–9, 12–14]. In particular, these magnetically charged black holes are known to be characterized by *infinite* spectra of imaginary (unstable) resonant frequencies $\{\omega_n(r_+, r_-)\}_{n=0}^{n=\infty}$. Based on direct *numerical* computations of the black-hole instability spectra [14], it has recently been pointed out [20] that the excited instability eigenvalues of the magnetically charged SU(2) Reissner-Nordström black holes are described, to a very good degree of accuracy, by the simple universal relation (1.3).

In the present paper we have studied analytically the characteristic instability spectra of the magnetically charged SU(2) Reissner-Nordström black-hole spacetimes in the small frequency regime. In particular, we have provided a simple *analytical* proof for the *numerically*-observed [14, 20] universal behavior [see eq. (1.3)]

$$\omega_n(r_+ - r_-) = \lambda_n \quad \text{for} \quad \omega_n r_+ \ll (r_+ - r_-)/r_+ \quad (6.1)$$

which characterizes the black-hole excited instability resonances, where $\{\lambda_n\}$ are constants. Our analysis has revealed that these dimensionless constants (which are *independent* of the black-hole parameters) are given by the simple relation [see eqs. (4.3) and (4.4)]

$$\lambda_n = i \times 8e^{-\frac{2\pi}{\sqrt{3}}(n-\frac{1}{3})+\frac{4\theta-2\phi}{\sqrt{3}}}. \quad (6.2)$$

Finally, it is interesting to note that one finds from (4.3) the simple dimensionless ratio

$$\frac{\omega_{n+1}}{\omega_n} = e^{-2\pi/\sqrt{3}} \quad (6.3)$$

for the characteristic excited instability eigenvalues of the magnetically charged SU(2) Reissner-Nordström black holes. It is worth emphasizing the fact that the relation (6.3), which characterizes the instability resonance spectra of the magnetically charged black-hole spacetimes, is *universal* in the sense that it is *independent* of the physical parameters (masses and magnetic charges) of the SU(2) Reissner-Nordström black holes.

Acknowledgments

This research is supported by the Carmel Science Foundation. I would like to thank Oliver Rinne for sharing with me his numerical data. I would also like to thank Yael Oren, Arbel M. Ongo, Ayelet B. Lata, and Alona B. Tea for helpful discussions.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] V. Moncrief, *Odd-parity stability of a Reissner-Nordström black hole*, *Phys. Rev. D* **9** (1974) 2707 [[INSPIRE](#)].
- [2] V. Moncrief, *Stability of Reissner-Nordström black holes*, *Phys. Rev. D* **10** (1974) 1057 [[INSPIRE](#)].
- [3] S. Hod, *Stability of the extremal Reissner-Nordström black hole to charged scalar perturbations*, *Phys. Lett. B* **713** (2012) 505 [[arXiv:1304.6474](#)] [[INSPIRE](#)].
- [4] S. Hod, *No-bomb theorem for charged Reissner-Nordström black holes*, *Phys. Lett. B* **718** (2013) 1489 [[INSPIRE](#)].
- [5] S. Hod, *Stability of highly-charged Reissner-Nordström black holes to charged scalar perturbations*, *Phys. Rev. D* **91** (2015) 044047 [[arXiv:1504.00009](#)] [[INSPIRE](#)].
- [6] P.B. Yasskin, *Solutions for gravity coupled to massless gauge fields*, *Phys. Rev. D* **12** (1975) 2212 [[INSPIRE](#)].
- [7] N. Straumann and Z.-H. Zhou, *Instability of the Bartnik-McKinnon solution of the Einstein Yang-Mills equations*, *Phys. Lett. B* **237** (1990) 353 [[INSPIRE](#)].
- [8] N. Straumann and Z.H. Zhou, *Instability of a colored black hole solution*, *Phys. Lett. B* **243** (1990) 33 [[INSPIRE](#)].
- [9] P. Bizoń and R.M. Wald, *The $n = 1$ colored black hole is unstable*, *Phys. Lett. B* **267** (1991) 173 [[INSPIRE](#)].
- [10] P. Breitenlohner, P. Forgács and D. Maison, *Gravitating monopole solutions*, *Nucl. Phys. B* **383** (1992) 357 [[INSPIRE](#)].
- [11] P. Breitenlohner, P. Forgács and D. Maison, *Gravitating monopole solutions II*, *Nucl. Phys. B* **442** (1995) 126 [[gr-qc/9412039](#)] [[INSPIRE](#)].
- [12] S. Hod, *The instability spectrum of weakly-magnetized SU(2) Reissner-Nordström black holes*, *Phys. Lett. B* **739** (2014) 157 [[arXiv:1410.7406](#)] [[INSPIRE](#)].
- [13] S. Hod, *Lifetime of unstable hairy black holes*, *Phys. Lett. B* **661** (2008) 175 [[arXiv:0803.0608](#)] [[INSPIRE](#)].
- [14] O. Rinne, *Formation and decay of Einstein-Yang-Mills black holes*, *Phys. Rev. D* **90** (2014) 124084 [[arXiv:1409.6173](#)] [[INSPIRE](#)].
- [15] C. Gundlach and J.M. Martín-García, *Critical phenomena in gravitational collapse*, *Living Rev. Rel.* **10** (2007) 5 [[arXiv:0711.4620](#)] [[INSPIRE](#)].
- [16] M.W. Choptuik, T. Chmaj and P. Bizoń, *Critical behavior in gravitational collapse of a Yang-Mills field*, *Phys. Rev. Lett.* **77** (1996) 424 [[gr-qc/9603051](#)] [[INSPIRE](#)].
- [17] C. Gundlach, *Echoing and scaling in Einstein-Yang-Mills critical collapse*, *Phys. Rev. D* **55** (1997) 6002 [[gr-qc/9610069](#)] [[INSPIRE](#)].
- [18] M.W. Choptuik, E.W. Hirschmann and R.L. Marsa, *New critical behavior in Einstein-Yang-Mills collapse*, *Phys. Rev. D* **60** (1999) 124011 [[gr-qc/9903081](#)] [[INSPIRE](#)].
- [19] P. Bizoń and T. Chmaj, *Remark on formation of colored black holes via fine tuning*, *Phys. Rev. D* **61** (2000) 067501 [[gr-qc/9906070](#)] [[INSPIRE](#)].

- [20] S. Hod, *Numerical evidence for universality in the excited instability spectrum of magnetically charged Reissner-Nordström black holes*, *Eur. Phys. J. C* **75** (2015) 180 [[arXiv:1504.04010](#)] [[INSPIRE](#)].
- [21] P. Bizoń, *Stability of Einstein Yang-Mills black holes*, *Phys. Lett. B* **259** (1991) 53 [[INSPIRE](#)].
- [22] S. Chandrasekhar, *On the equations governing the perturbations of the Schwarzschild black hole*, *Proc. Roy. Soc. Lond. A* **343** (1975) 289.
- [23] S.A. Teukolsky and W.H. Press, *Perturbations of a rotating black hole. III. Interaction of the hole with gravitational and electromagnetic radiation*, *Astrophys. J.* **193** (1974) 443 [[INSPIRE](#)].
- [24] D.N. Page, *Particle emission rates from a black hole: massless particles from an uncharged, nonrotating hole*, *Phys. Rev. D* **13** (1976) 198 [[INSPIRE](#)].
- [25] S. Hod, *Quasinormal resonances of near-extremal Kerr-Newman black holes*, *Phys. Lett. B* **666** (2008) 483 [[arXiv:0810.5419](#)] [[INSPIRE](#)].
- [26] S. Hod, *Purely imaginary polar resonances of rapidly-rotating Kerr black holes*, *Phys. Rev. D* **88** (2013) 084018 [[arXiv:1311.3007](#)] [[INSPIRE](#)].
- [27] M. Abramowitz and I.A. Stegun, *Handbook of mathematical functions*, Dover Publications, New York U.S.A. (1970).